

ON THE HOMOTOPY OF FINITE CW-COMPLEXES WITH POLYCYCLIC FUNDAMENTAL GROUP

Mihai DAMIAN
 Université Louis Pasteur
 IRMA, 7, rue René Descartes,
 67 084 STRASBOURG
 e-mail : damian@math.u-strasbg.fr
 june 2006

Abstract : Let X be a finite connected CW-complex of dimension q . If its fundamental group $\pi_1(X)$ is polycyclic of Hirsch number $h > q$, we show that at least one homotopy group $\pi_i(X)$ is not finitely generated. If $h = q$ or $h = q - 1$ the same conclusion holds unless X is an Eilenberg-McLane space $K(\pi_1(X), 1)$.

Mathematics subject classification : 57R70; 55P15; 57Q10.

Keywords : Novikov homology, closed 1-forms, CW-complexes, homotopy groups.

1 Introduction

Let X be a finite connected CW-complex of dimension q . Consider the homotopy groups $\pi_i(X) = [\mathbf{S}^i, X]$, for $i \geq 2$. If all these (Abelian) groups are finitely generated we say that the homotopy of X is finitely generated. For simply connected complexes, a celebrated theorem of Serre [30] asserts that

Theorem 1.1 *If $\pi_1(X) = 1$ the homotopy of X is finitely generated.*

On the other hand simple examples as $X = \mathbf{S}^1 \vee \mathbf{S}^2$ show that the homotopy of X is not always finitely generated.

One can then ask if there is a general negative statement asserting that the homotopy of X is not finitely generated under some hypothesis on computable invariants of X , such as its fundamental group. This is the aim of the present paper. The statements we will prove have the following form :

If the fundamental group of X satisfies to the conditions (C) (which depend on the dimension q), then the homotopy of X is not finitely generated unless X is an Eilenberg-McLane space $K(\pi_1(X))$.

If X is an Eilenberg-McLane space, the cohomological dimension of $\pi_1(X)$ is less or equal than $q = \dim(X)$. So, we may add $cd(\pi_1(X)) > q$ to the hypothesis (C) in order to have the desired conclusion on the homotopy of X . For instance when $\pi_1(X)$ has non trivial torsion elements then $cd(\pi_1(X))$ is infinite and the above conclusion is valid.

1.1 Statement of the results

Before stating the main theorem we remind the following :

Definition. A group G is called polycyclic if it admits a series

$$1 = G_0 < G_1 < \cdots < G_k = G$$

with cyclic factors G_{i+1}/G_i . When all the factor groups are infinite cyclic we call G a poly- $\mathbf{Z}$ group.

K.A. Hirsch proved in [14] that the number of infinite cyclic factors in such a series is an invariant of G . It is called Hirsch number of G and it is denoted by $h(G)$.

Here are our main statements :

Theorem 1.2 *Suppose that X is a finite connected CW complex of dimension q and that $\pi_1(X)$ is polycyclic. Then*

- a) If $h(\pi_1(X)) > q$, the homotopy of X is not finitely generated.*
- b) If $h(\pi_1(X)) \in \{q-1, q\}$ then the homotopy of X is not finitely generated unless X is a $K(\pi_1(X), 1)$. In particular, when $\pi_1(X)$ has torsion then the conclusion of a) holds.*

Note that, by a result of J-P. Serre (see 4.2 below), when the homotopy of X is not finitely generated at least one of the groups $\pi_2(X), \pi_3(X), \dots, \pi_q(X)$ is not finitely generated. When X is a manifold we can improve this result :

Theorem 1.3 *Let M^n be a closed manifold. Let $r = \max \left\{ \left\lfloor \frac{n}{2} \right\rfloor, 3 \right\}$. Suppose that $\pi_1(M)$ is polycyclic. Then, if $h(\pi_1(M)) \geq n - 1$, the group $\pi_i(M)$ is not finitely generated for some $i \leq r$ unless $h(\pi_1(M)) = n$ and M is a $K(\pi_1, 1)$.*

Moreover, if $n \geq 6$ we can replace the assertion " M is a $K(\pi_1, 1)$ " by the stronger one "the universal cover of M is diffeomorphic to $\mathbf{R}^n$ ".

When $n = 4$, we are able to prove the above statement for $r = 2$, namely :

Theorem 1.4 *Let M^4 be a closed connected manifold with polycyclic fundamental group. If $h(\pi_1(M)) \geq 3$ then $\pi_2(M)$ is not finitely generated unless $h(\pi_1(M)) = 4$ and M is a $K(\pi_1, 1)$.*

In the next theorem we weaken the hypothesis on the fundamental group but we suppose in addition that $\chi(X) \neq 0$. Recall first the finiteness properties of a group, which were introduced by C.T.C. Wall in [35] :

Definition. Let $r \geq 1$ be an integer. A groupe G is of type $\mathcal{F}_r$ if there is an Eilenberg-McLane space $K(G, 1)$ whose r -skeleton has a finite number of cells. Equivalently a group is of type $\mathcal{F}_r$ if it acts freely, properly, cellularly and cocompactly on an $(r - 1)$ -connected cell complex.

A group is of type $\mathcal{F}_\infty$ if it is of type $\mathcal{F}_r$ for any integer $r > 0$.

Remark. A group G is of type $\mathcal{F}_1$ if and only if it is finitely generated. G is of type $\mathcal{F}_2$ if and only if it is finitely presented.

Theorem 1.5 *Let X^q be a finite connected CW-complex of dimension q with fundamental group of type $\mathcal{F}_{q+1}$. Suppose that there is a non vanishing morphism $u : \pi_1(X) \rightarrow \mathbf{Z}$ such that $\text{Ker}(u)$ is of type $\mathcal{F}_{q+1}$. Then, if $\chi(X) \neq 0$, the homotopy of X is not finitely generated.*

If X is a manifold with non-zero Euler characteristic we obtain the same holds for $\pi_1(X)$ and $\text{Ker}(u)$ of type $\mathcal{F}_{\lfloor \frac{q}{2} \rfloor}$.

The hypothesis on u may be reformulated in terms of the Bieri-Renz invariants of $\pi_1(X)$ as follows :

$$(*) \quad \Sigma^{q+1}(\pi_1) \cap -\Sigma^{q+1}(\pi_1) \neq \emptyset.$$

The Bieri-Renz invariants $\Sigma^i(G)$ of a group G are open subsets of the unit sphere of $\mathbf{R}^{rk(G)}$. We recall the definition and the properties of these invariants in Section 4.

In the next subsection we show that polycyclic fundamental groups satisfy the hypothesis of 1.5 and we give other examples of groups for which the condition $(*)$ is fulfilled.

1.2 Comments on the results

An Abelian group is obviously polycyclic of Hirsch number equal to its rank. It is also of type $\mathcal{F}_\infty$. For Abelian fundamental groups the results of the previous subsection were proved by the author in [9]. Here are other remarks about these statements.

Remarks. 1. The lower bound $q - 1$ for $h(\pi_1)$ in the hypothesis of 1.2 is optimal. Indeed, the complex $X = \mathbf{T}^{n-2} \times \mathbf{S}^2$ fulfills the conditions of the hypothesis of 1.2 except for $h(\pi_1(X)) = q - 2$. It is a consequence of the theorem 1.1 of J.P.Serre, quoted above, that X does not satisfy to the conclusion of 1.2.

2. If G is a polycyclic group then $[G, G]$ is obviously polycyclic and therefore for every morphism $u : G \rightarrow \mathbf{Z}$, $\text{Ker}(u)$ has the same property. This implies that the hypothesis on the fundamental group of 1.5 is fulfilled when $\pi_1(X)$ is polycyclic. Indeed we have :

Proposition 1.6 *Polycyclic groups are of type $\mathcal{F}_\infty$.*

Proof

The proposition is an immediate corollary of the following :

Lemma 1.7 *Consider an exact sequence of groups*

$$1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1.$$

If K and H are of type $\mathcal{F}_\infty$ then so is G .

To prove the above lemma one may use the following results :

- a) G is $\mathcal{F}_\infty$ iff $H_*(G, \cdot)$ commutes with direct products. This was proved by R. Bieri and B. Eckmann in [2].
- b) The Hochschild-Serre spectral sequence [16] which satisfies

$$E_{pq}^2 = H_p(Q, H_q(K, R))$$

and converges towards $H_*(G, R)$. (The complete statement 4.6 is given in Section 4).

It is clear how a) and b) imply 1.7 proving thus 1.6. $\diamond$

3. Other examples of groups satisfying the hypothesis of 1.5 were constructed by M. Bestvina, N. Brady in [1] and then in a more general statement by J. Meier, H. Meinert and L. Van Wyk in [20]. Let us describe them briefly :

Definition. A flag complex L is a finite simplicial complex with the property that any collection of $q + 1$ mutually adjacent vertices span a q -simplex in L . The *right angled Artin group* associated to L is the group G_L spanned by the vertices $v_1, \dots, v_s$ of L with relations $[v_i, v_j] = 1$ whenever v_i and v_j are adjacent in L .

Note that the group G_L admits a finite Eilenberg-McLane space [1] and therefore it is of type $\mathcal{F}_\infty$. We have

Theorem 1.8 (*Bestvina, Brady, Meier, Meinert, van Wyk*) *Let L and G_L as above and consider a morphism $u : G_L \rightarrow \mathbf{Z}$ such that $u(v_i) \neq 0$ for each generator v_i . Then, if L is $(r - 1)$ connected, $\text{Ker}(u)$ is of type $\mathcal{F}_r$.*

Considering appropriate flag complexes L (remark that the barycentric subdivision of any complex is a flag complex) we may find a lot of examples of groups which fulfil the hypothesis of 1.5.

1.3 Idea of the proof

The general idea of the proof of 1.3 is the following : we show first that a manifold M as in 1.3 whose dimension is greater or equal to 6 and whose homotopy groups $\pi_i(M)$ are of finite type for $i \leq r$ admits a fibration over the circle. This is the main difficulty of the proof. To overcome it, we will use

Novikov homology theory and some of its applications on Morse functions $f : M \rightarrow \mathbf{S}^1$.

Remark that if the manifold F^{n-1} is a fiber, the inclusion $j : F \hookrightarrow M$ induces isomorphisms in π_i for $i \geq 2$. The lift of j induces a homotopy equivalence between the universal covers of F and M .

Now take a manifold M , as in 1.3 and consider the product $M \times \mathbf{S}^3$ in order to fulfill the condition on the dimension. Supposing that the homotopy groups $\pi_i(M)$ are of finite type for $i \leq r$, apply the previous argument to this product and get a fiber F as above. Then check that F still satisfies the hypothesis of 1.3. If its dimension is still greater or equal than 6 we apply the same argument to F . By successive iterations we find a manifold F_0 of low dimension whose universal cover has the same homotopy type as the one of $M \times \mathbf{S}^3$. In particular the homology groups of $\tilde{F}_0$ and $\tilde{M} \times \mathbf{S}^3$ are isomorphic. By comparing them, we infer that the universal cover of M is acyclic, which means that M is Eilenberg-McLane. So, either the homotopy of M is not finitely generated, or M is Eilenberg-McLane, as in the statement of 1.3.

To prove 1.2 we embed X into an Euclidian space and thicken it to a manifold W , with boundary M . Then we apply the above argument to M .

In the hypothesis of 1.5, supposing that the homotopy of X is finitely generated, we are only able to prove that *the Novikov homology* $H_*(M, u)$ of the corresponding manifold M vanishes. But this implies that $\chi(M) = \chi(X) = 0$, yielding a contradiction.

The paper is organized as follows. In Section 2 we state the result 2.1 on the fibration over the circle which was roughly sketched above. Supposing 2.1 true, we show how it can be successively applied in order to prove 1.3, 1.4 and 1.2. In Section 3 we recall the definition and some useful properties of the Novikov homology. We also recall some basic facts about Morse theory of circle-valued functions and point out the relation between the (vanishing of the) Novikov homology and the existence of a fibration over the circle. In Section 4 we prove 2.1 and 1.5. We will use the Bieri-Renz criterion which we recall in the Subsection 4.3, devoted to the Bieri-Renz invariants.

2 Iterated fibrations

Our main result 1.2 is a consequence of the following :

Theorem 2.1 *Let M^n a closed manifold of dimension $n \geq 6$. Suppose that π_1 is of type $\mathcal{F}_{[\frac{n}{2}]}$. Suppose also that $\pi_i(M)$ are of finite type for $i \leq [\frac{n}{2}]$.*

Suppose that there is a non zero cohomology class $u \in H^1(M; \mathbf{R}) \approx \text{Hom}(\pi_1(M), \mathbf{R})$ such that $\text{Ker}(u)$ is of type $\mathcal{F}_{[\frac{n}{2}]}$. Suppose also that the Whitehead group $Wh(\pi_1(M))$ vanishes.

*Then there is a fibration $f : M \rightarrow \mathbf{S}^1$ such that $[f^*d\theta] = u$.*

When M of arbitrary dimension n and $\text{Ker}(u)$ is of type $\mathcal{F}_r$, where $r = \max\{[\frac{n}{2}], 2\}$ the same conclusion holds if we replace M by $M \times \mathbf{S}^p$ for all $p \geq 6 - n$.

Remarks. 1. The Whitehead group is defined as follows :

$$Wh(\pi) = \frac{GL\mathbf{Z}[\pi]}{[GL\mathbf{Z}[\pi], GL\mathbf{Z}[\pi]], \{\pm g | g \in \pi\}},$$

where $GL\mathbf{Z}[\pi] = \lim_{\rightarrow} GL_m\mathbf{Z}[\pi]$.

2. T. Farrell and W. Hsiang proved in [11] that $Wh(\pi)$ vanishes when π is poly- $\mathbf{Z}$. The Whitehead group also vanishes when $\pi = G_L$ is one of the examples of Bestvina and Brady. This result was proved by B. Hu [17].

It is conjectured that $Wh(\pi) = 0$ for any torsion-free group π .

3. For $\pi_1 = \mathbf{Z}$, Theorem 2.1 was proved by W. Browder and J. Levine in [6].

The proof of 2.1 will be given in Section 4. Let us now show how this theorem implies our main results 1.3, 1.4 and 1.2.

Proof of 2.1 $\implies$ 1.3

Without restricting the generality of our statements, we may suppose that $n \geq 3$. We begin with the statement of the following result, due to K. Hirsch ([15], Theorem 2).

Theorem 2.2 *Let G be a polycyclic group. There exists a normal subgroup N of finite index in G which is poly- $\mathbf{Z}$ and such that $h(N) = h(G)$.*

Now, let $\pi_1^0 \leq \pi_1$ a subgroup as in 2.2 and consider the associated finite cover $M_0 \rightarrow M$. Let

$$1 = G_0 < G_1 < \cdots < G_k = \pi_1^0$$

be a series with infinite cyclic factor groups : we have therefore $h(\pi_1) = k$, so by hypothesis $k \geq n-1$. Denote by u_1 the projection $G_k \rightarrow G_k/G_{k-1} \approx \mathbf{Z}$.

Suppose that $\pi_i(M)$ is finitely generated for $i \leq r = \max \left\{ \left\lfloor \frac{n}{2} \right\rfloor, 3 \right\}$.

Consider first the case $n \geq 6$. Using 1.6 we find that u_1 fulfills the hypothesis of 2.1. We apply 2.1 and we get a fibration $M_0 \rightarrow \mathbf{S}^1$. Let F_1 be a fibre of this fibration. So F_1 satisfies :

1. $\pi_1(F_1) = \text{Ker}(u_1) = G_{k-1}$.
2. $\pi_i(F_1) \approx \pi_i(M) \forall i \geq 2$.

We wish to apply 2.1 to F_1 . We use the cohomology class $u_2 : G_{k-1} \rightarrow G_{k-1}/G_{k-2}$. Its kernel is G_{k-2} , so, using again 1.6, $\text{Ker}(u_2)$ is of type $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor}$ (actually $\mathcal{F}_\infty$, since polycyclic). The hypothesis on the higher homotopy groups of 2.1 is fulfilled by F_1 (because of the condition 2. above), therefore we may apply 2.1 to F_1 if its dimension is no less than 6.

We get thus a closed connected manifold $F_2 \subset F_1$ whose higher homotopy groups are those of M and whose fundamental group is G_{k-2} . If its dimension is greater or equal than 6 we may again apply 2.1 to the couple $(F_2, u_3 : G_{k-2} \rightarrow G_{k-2}/G_{k-3})$.

By iterating this argument we get a sequence

$$(1) \quad F_{n-5} \hookrightarrow F_{n-6} \hookrightarrow \dots \hookrightarrow F_1 \hookrightarrow M_0$$

such that for $j = 1, \dots, n-1$:

1. $\pi_1(F_j) = G_{k-j}$
2. $\dim(F_j) = n-j$.
3. The inclusion $F_j \hookrightarrow M$ induces isomorphisms in π_i for $i \geq 2$.

The manifold F_{n-5} is of dimension 5, so it does not verify the dimension hypothesis of 2.1. Its fundamental group is G_{k-n+5} . In order to continue to apply 2.1, we consider the product $F = F_{n-5} \times \mathbf{S}^3$. As above, we have the cohomology class $u_{k-n+4} : G_{k-n+5} \rightarrow G_{k-n+5}/G_{k-n+4}$ which is non zero and has a (poly- $\mathbf{Z}$) kernel of type $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor}$. By 2.1, we get a fibration of F over $\mathbf{S}^1$. Its fiber K_0 is a closed connected manifold of dimension 7 with finitely generated π_i for $i = 1, \dots, r = \max \left\{ \left\lfloor \frac{n}{2} \right\rfloor, 3 \right\}$ (since $\mathbf{S}^3$ has the same property by 1.1). Its fundamental group is G_{k-n+4} . Since $\lfloor 7/2 \rfloor = 3$, we may apply 2.1 to K_0 and then again to its submanifold K_1 given by 2.1, to obtain a sequence as above, where the maps are inclusions of fibers of fibrations over

the circle which therefore induce isomorphisms at the level of π_i for $i \geq 2$:

$$K_2 \hookrightarrow K_1 \hookrightarrow K_0 \hookrightarrow F.$$

It follows that the universal covers of K_2 and F are homotopically equivalent, in particular

$$(2) \quad H_*(\widetilde{K}_2) \approx H_*(\widetilde{F}_{n-5} \times \mathbf{S}^3).$$

Now K_2 is a closed connected 5-dimensional manifold whose fundamental group is $G = G_{k-n+2}$. Since $k \geq n - 1$, G is an infinite group. It follows that $H_5(\widetilde{K}_2) = 0$ (see [5], p.346) ; We have therefore that $H_i(\widetilde{K}_2)$ vanishes for $i > 4$.

Using the Kunneth formula we infer from (2) that $H_i(\widetilde{F}_{n-5})$ vanishes for $i > 0$. Therefore $\widetilde{F}_{n-5}$ is contractible and, using the sequence (1) $\widetilde{M}_0 = \widetilde{M}$ is contractible, too. So M is an Eilenberg-Mc Lane space $K(\pi_1(M), 1)$. The cohomological dimension of $\pi_1(M)$ is therefore equal to n . Now we use the following well-known result (for a proof, see [12], Lemma 8, p. 154) :

Proposition 2.3 *If G is a polycyclic group then $cd(G)$ is finite if and only if G is torsion free. In this case $cd(G) = h(G)$.*

Applying 2.3 we get that M cannot be a $K(\pi_1, 1)$ unless $h(\pi_1(M)) = n$. If this relation is not valid, we get a contradiction and therefore the homotopy of M cannot be finitely generated, completing the proof of 1.3 for $n \geq 6$. If it is and if M is a $K(\pi_1, 1)$, let us show that its universal cover is $\mathbf{R}^n$. Since according to 2.1 M fibers over $\mathbf{S}^1$, we have

$$\widetilde{M} \approx \widetilde{F} \times \mathbf{R}.$$

To finish the proof we just have to apply a celebrated theorem of J. Stallings [29] which asserts :

Theorem 2.4 *If a manifold $P^{n \geq 5}$ is a product of two open non-trivial contractible manifolds then P is diffeomorphic to $\mathbf{R}^n$.*

◇

Proof of 2.1 $\implies$ 1.4 (case $n=4$)

Suppose that $\pi_2(M)$ is finitely generated. As in the proof of 1.3, use 2.2 and consider a finite cover M_0 with poly- $\mathbf{Z}$ fundamental group π_1^0 of Hirsch number greater than 3. We apply first 2.1 to $M_0 \times \mathbf{S}^2$ and to the morphism $u_1 : G_k \rightarrow G_k/G_{k-1}$. We get a manifold F of dimension 5 which is a fiber of a fibration $M_0 \rightarrow \mathbf{S}^1$. Then we apply again 2.1 to the product $F \times \mathbf{S}^2$ and the cohomology class $u_2 : G_{k-1} \rightarrow G_{k-1}/G_{k-2}$ and we get a 6-dimensional submanifold $K \hookrightarrow F \times \mathbf{S}^2$. We have a sequence

$$K \hookrightarrow F \times \mathbf{S}^2 \hookrightarrow M \times \mathbf{S}^2 \times \mathbf{S}^2,$$

which induces homotopic equivalences at the level of the universal covers. In particular

$$H_*(\widetilde{K}) \sim H_*(\widetilde{M} \times \mathbf{S}^2 \times \mathbf{S}^2).$$

Now, as above, since $k \geq 3$, the fundamental group G_{k-2} of K is infinite, so the homology of $\widetilde{K}$ vanishes in degrees $i > 5$. This implies that the homology of $\widetilde{M}$ is zero, so M is a $K(\pi_1, 1)$. We conclude using 2.3. $\diamond$

Proof of 2.1 $\implies$ 1.3 in the cases $n = 3, 5$

Suppose that $\pi_2(M)$ and $\pi_3(M)$ are finitely generated. We apply $(n - 2)$ times Theorem 2.1 to the 8-dimensional manifold $M \times \mathbf{S}^{8-n}$. We get a sequence :

$$K_{n-2} \hookrightarrow K_{n-3} \cdots \hookrightarrow M \times \mathbf{S}^{8-n},$$

as above. The manifold K_{n-2} is of dimension $8 - n + 2$ and has an infinite fundamental group. So, $H_i(\widetilde{K}_{n-2}) = 0$ for $i \geq 8 - n + 2$. Since this homology is isomorphic to $H_*(\widetilde{M} \times \mathbf{S}^{8-n})$, it follows that $\widetilde{M}$ is acyclic, and therefore that M is an Eilenberg-McLane. As above, we use 2.3 to finish the proof. $\diamond$

Proof of 2.1 $\implies$ 1.2

Assume that the homotopy of X is finitely generated.

a) Suppose that $h(\pi_1(X)) = k$ for some integer $k > q$. Using 2.2, consider a finite cover X_0 of X such that $\pi_1(X_0)$ is poly- $\mathbf{Z}$ and $h(\pi_1(X_0)) = k$.

Embed X_0 in an Euclidian space $\mathbf{R}^{2q+r+1}$ for some $r \geq 0$ which will be fixed later in the proof. Let W be a tubular neighbourhood of X_0 and denote by M^{2q+r} the smooth manifold ∂W . Since M is a deformation retract of $W \setminus X_0$, using a general position argument we get isomorphisms between

$\pi_i(W)$ and $\pi_i(M)$ for $i \leq q + r - 1$. In particular, for $r > 1$ the inclusion $M \hookrightarrow W$ induces an isomorphism at the level of fundamental groups and, since X_0 is a retract of W , the higher homotopy groups $\pi_i(M)$ are finitely generated for $i \leq q + r - 1$. If $r \geq 1$ then $q + r - 1 \geq \left\lfloor \frac{2q+r}{2} \right\rfloor$ and the hypothesis on the higher homotopy groups of 2.1 is satisfied by the manifold M . Let

$$1 = G_0 < G_1 < \cdots < G_k = \pi_1(X_0)$$

be a series with infinite cyclic factor groups. For r large enough ($r \geq k - 2q + 5$), we may apply k times 2.1, as in the proof of 1.3, and get a sequence of closed connected manifolds :

$$(3) \quad F_k \hookrightarrow F_{k-1} \hookrightarrow \cdots \hookrightarrow F_1 \hookrightarrow M$$

such that for $j = 0, \dots, k$:

1. $\pi_1(F_j) = G_{k-j}$.
2. $\dim(F_j) = 2q + r - j$.
3. The inclusion $F_j \hookrightarrow M$ induces isomorphisms in π_i for $i \geq 2$.

The constant $r \geq 1$ must be chosen large enough to have :

- a) $\dim(F_j) \geq 5 \ \forall j$: needed for the dimension hypothesis in 2.1 (this means $2q + r - k \geq 5$).
- b) $r + q - 1 \geq \left\lfloor \frac{2q+r}{2} \right\rfloor$, (which is true for $r \geq 1$) to insure the hypothesis on the higher homotopy groups in 2.1, as we explained above.
- c) $r > k - q$.

It follows that the first manifold F_k in the sequence (1) is closed and simply connected, of dimension $2q + r - k > q$. Using the property 3 we find that the composition

$$(4) \quad F_k \hookrightarrow M \hookrightarrow W \rightarrow X_0$$

induces isomorphisms at the level of the higher homotopy groups π_i for $i = 2, \dots, q + r - 1$. By Whitehead's theorem, the induced application between the universal covers of F_k and X_0 is an isomorphism at the level of the i^{th} homology group for $i \leq q + r - 1$. In particular, since $q + r - 1 \geq 2q + r - k$ (since by hypothesis $k > q$), we have

$$(5) \quad H_{2q+r-k}(\tilde{F}_k) \approx H_{2q+r-k}(\tilde{X}_0).$$

But F_k is simply connected of dimension $2q + r - k$, so the left side of (5) is $\mathbf{Z}$. On the other hand $2q + r - k > q = \dim(X_0)$, so the right side of (5) vanishes. This contradicts the initial assumption on the finite generation of the homotopy of X and the statement a) of the theorem is proved.

b) Suppose now that $k = h(\pi_1(X)) \in \{q-1, q\}$. As above we get a closed simply connected manifold F_k of dimension $2q+r-k$ (i.e. $q+r+1$ or $q+r$) such that the application $H_i(F_k) \rightarrow H_i(\widetilde{X}_0)$ is an isomorphism for $i \leq q+r-1$. In particular, the homology $H_i(F_k; \mathbf{Z})$ vanishes for $i = q+1, \dots, q+r-1$.

By the univesal coefficients theorem we obtain that the cohomology $H^i(F_k; \mathbf{Z})$ is zero for $i = q+2, \dots, q+r-1$, so Poincaré duality implies that $H_i(F_k; \mathbf{Z})$ also vanishes for $i = q-k+1, \dots, q+r-k-2$. Now $q-k+1 \in \{1, 2\}$ and F_k is simply connected : We infer after choosing r sufficiently large that the integer homology of F_k vanishes in the degrees $i \leq q$ which means that $\widetilde{X}_0$ is contractible. Therefore X_0 and X are Eilenberg-Mac Lane spaces. In particular the cohomological dimension of $\pi_1(X)$ cannot exceed q , which implies that $\pi_1(X)$ is torsion free.

This completes the proof. $\diamond$

3 Novikov homology and fibrations over the circle

In the preceeding section we showed that our main results 1.2, 1.3 and 1.4 are consequences of 2.1. Let M be a closed connected manifold and $u \in H^1(M; \mathbf{Z}) \approx \text{Hom}(\pi_1, \mathbf{Z})$. The aim of the present section is to introduce the Novikov homology $H_*(M, u)$ and to describe the situations when the vanishing of $H_*(M, u)$ implies the existence of a fibration $f : M \rightarrow \mathbf{S}^1$ i.e. the conclusion of 2.1. Then in the Section 4 we prove that the hypothesis of 2.1 implies the vanishing of the Novikov homology $H_*(M, u)$.

3.1 Novikov homology

Let $u \in H^1(M; \mathbf{R})$. Denote by Λ the ring $\mathbf{Z}[\pi_1(M)]$ and by $\hat{\Lambda}$ the ring of formal series $\mathbf{Z}[[\pi_1(M)]]$. Consider a $\mathcal{C}^1$ -triangulation of M which we lift to the universal cover $\widetilde{M}$. We get a Λ -free complex $C_\bullet(M)$ spanned by (fixed lifts of) the cells of the triangulation of M .

We define now *the completed ring* Λ_u :

$$\Lambda_u := \left\{ \lambda = \sum n_i g_i \in \hat{\Lambda} \mid g_i \in \pi_1(M), n_i \in \mathbf{Z}, u(g_i) \rightarrow +\infty \right\}$$

The convergence to $+\infty$ means here that for all $A > 0$, $u(g_i) < A$ only for a finite number of g_i which appear with a non-zero coefficient in the sum λ .

Remark Let $\lambda = 1 + \sum n_i g_i$ where $u(g_i) > 0$ for all i . Then λ is invertible in Λ_u . Indeed, if we denote by $\lambda_0 = \sum n_i g_i$ then it is easy to check that $\sum_{k \geq 0} (-\lambda_0)^k$ is an element of Λ_u and it is obvious that it is the inverse of λ .

Definition Let $C_\bullet(M, u)$ be the Λ_u -free complex $\Lambda_u \otimes_\Lambda C_\bullet(M)$. The Novikov homology $H_*(M, u)$ is the homology of the complex $C_\bullet(M, u)$.

A purely algebraic consequence of the previous definition is the following version of the universal coefficients theorem ([13], p.102, Th 5.5.1) :

Theorem 3.1 *There is a spectral sequence E_{pq}^r which converges to $H_*(M, u)$ and such that*

$$E_{pq}^2 = \text{Tor}_p^\Lambda(H_q(\widetilde{M}), \Lambda_u).$$

We will use this result in Section 4 to prove that in the hypothesis of 1.3, the Novikov homology associated to some class vanishes.

3.2 Morse-Novikov theory

We recall in this subsection the relation between Novikov homology and closed one forms. In dimension $n \geq 6$, when the Novikov homology vanishes, some hypothesis on π_1 stated below imply the existence of a nowhere vanishing closed one form on M . It is well-known (see [34]) that the existence of such a form is equivalent to the existence of a fibration of M over $\mathbf{S}^1$.

Let α be a closed generic one form in the class u . Let ξ be the gradient of α with respect to some generic metric on M . For every critical point c of α we fix a point $\tilde{c}$ above c in the universal cover $\widetilde{M}$. We can define then a complex $C_\bullet(\alpha, \xi)$ spanned by the zeros of α : the incidence number $[d, c]$ for two zeros of consecutive indices is the (possibly infinite) sum $\sum n_i g_i$ where n_i is the algebraic number of flow lines which join c and d and which are covered by a path in $\widetilde{M}$ joining $g_i \tilde{c}$ and $\tilde{d}$. It turns out that this incidence number belongs to Λ_u , so $C_\bullet(\alpha, \xi)$ is actually a Λ_u -free complex.

The fundamental property of the Novikov homology is that it is isomorphic to the homology of the complex $C_\bullet(\alpha, \xi)$ above for any couple (α, ξ) .

By comparing the complexes $C_\bullet(\alpha, \xi)$ and $C_\bullet(-\alpha, -\xi)$ we get the following duality property (see Prop.2.8 in [8] and 2.30 in [18]) :

Theorem 3.2 *Let M^n be a closed connected manifold, $u \in H^1(M; \mathbf{R})$ and let l be an integer. If $H_i(M, -u) = 0$ for $i \leq l$, then $H_i(M, u) = 0$ for $i \geq n - l$.*

If the form α has no zeroes then $C_\bullet(\alpha, \xi)$ vanishes and therefore we have $H_*(M, [\alpha]) = 0$. Conversely, one can ask if the vanishing of $H_*(M, u)$ implies the existence of a nowhere vanishing closed 1-form belonging to the class $u \in H^1(M)$. For $n \geq 6$ this problem was independently solved by F. Latour [18] and A. Pajitnov [23], [24]. The statement is ([18], Th.1') :

Theorem 3.3 *For $\dim(M) \geq 6$ the following set of conditions is equivalent to the existence of a nowhere vanishing closed 1-form in $u \in H^1(M, \mathbf{Z})$:*

1. *Vanishing Novikov homology $H_*(M, u)$.*
2. *Vanishing Whitehead torsion $\tau(M, u) \in Wh(M, u)$.*
3. *Finitely presented $\text{Ker}(u) \subset \pi_1(M)$.*

Remarks

1. The definition of the generalized Whitehead group $Wh(M, u)$ and of the Whitehead torsion is given in [18].

2. In the statement of [23] the first two conditions are replaced by :

- 1'. $C_\bullet(M, u)$ is simply equivalent to zero.

Actually, one can show (see [19]) that 1' is equivalent to "1 and 2".

3. In earlier works on the subject as those of F.T. Farrell [10] and L. Siebenmann [31] the algebraic conditions which are equivalent to the existence of a nowhere vanishing closed 1-form in a rational cohomology class u were stated in the hypothesis that the infinite cyclic cover P associated to u is finitely dominated. ($P \rightarrow M$ is defined to be the pull-back of the universal covering $\mathbf{R} \rightarrow \mathbf{S}^1$ defined by a function $f : M \rightarrow \mathbf{S}^1$ such that $[f^*d\theta] = u$). The relation between the finite domination of P and vanishing of the Novikov homology was first established by A. Ranicki in [27], [28]. A Ranicki ([26], Chap.14) established a relation between the Whitehead "fiber obstruction" from [10] and [31] and the condition 2 above. Then A. Pajitnov and A. Ranicki proved in [25] the following

Theorem 3.4 *If G is a group with $Wh(G) = 0$ and $u : G \rightarrow \mathbf{Z}$ is a morphism, then $Wh(G, u) = 0$.*

It follows that under the hypothesis of 2.1 the condition 2 of 3.3 is always satisfied.

In order to prove 1.5 we use

Proposition 3.5 *If $H_*(M, u) = 0$ for some class u then $\chi(M) = 0$.*

Proof

The complex $C_\bullet(M, u)$ is acyclic. Like in [19] one can then show that this complex is simply equivalent to a complex of the form

$$0 \rightarrow \Lambda_u^p \xrightarrow{\partial} \Lambda_u^p \rightarrow 0.$$

This means that the second complex is isomorphic to the first after adding or cancelling a finite number of trivial summands

$$0 \rightarrow \Lambda_u \xrightarrow{Id} \Lambda_u \rightarrow 0.$$

In particular

$$\chi(M) = \chi(C_\bullet(M, u)) = 0.$$

◇

4 Novikov homology and finiteness properties of groups

In this section we will achieve the proof of our results 1.2 ... 1.5. Up to now we have proved that 1.2, 1.3 and 1.4 are implied by 2.1 (Section 2) and that the conclusion of 2.1 is valid if the three conditions of 3.3 are fulfilled. It remains to show that the hypothesis of 2.1 implies these three conditions. It is clearly the case for the third one, since in the hypothesis of 2.1, $Ker(u)$ is of type $\mathcal{F}_{[\frac{n}{2}]}$, in particular of type $\mathcal{F}_2$, i.e. finitely presented. The second condition is also satisfied, according to 3.4.

The main point of our proof is the vanishing of the Novikov homology $H_*(M, u)$ under the assumption of the hypothesis of 2.1.

Recall that, by 3.1, there is a spectral sequence E_{pq}^r which converges to $H_*(M, u)$ and whose term E_{pq}^2 is equal to $Tor_p^\Lambda(H_q(\widetilde{M}), \Lambda_u)$. Recall also that, by the duality result 3.2 we have the implication :

$$H_i(M, \pm u) = 0 \ \forall i \leq \left\lfloor \frac{n}{2} \right\rfloor \quad \Rightarrow \quad H_i(M, u) = 0 \ \forall i.$$

It suffices therefore to prove the following statement :

Proposition 4.1 *Let M^n a closed manifold. Suppose that $\pi_1(M)$ is of type $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor}$. Suppose also that $\pi_i(M)$ are of finite type for $i \leq \lfloor \frac{n}{2} \rfloor$.*

Suppose that there is a non zero cohomology class $u \in H^1(M; \mathbf{R}) \approx \text{Hom}(\pi_1(M), \mathbf{R})$ such that $\text{Ker}(u)$ is of type $\mathcal{F}_{\lfloor \frac{n}{2} \rfloor}$. Then, for all integers $0 \leq p, q \leq \lfloor \frac{n}{2} \rfloor$ we have

$$Tor_p^\Lambda(H_q(\widetilde{M}), \Lambda_{\pm u}) = 0.$$

Note that this result is purely algebraic. In order to proof it we use some facts about

4.1 Hurewicz-type morphisms

Recall that the classical Hurewicz's theorem asserts that for $q \geq 2$ the canonical morphism $I_q : \pi_q(M) \rightarrow H_q(M)$ is an isomorphism provided that M is $(q-1)$ -connected.

In [30] J-P. Serre generalized this theorem (see also [33], p. 504) : For some "admissible" classes of groups $\mathcal{C}$, he showed that, if X is simply connected such that $\pi_i(X) \in \mathcal{C}$ for $i = 1, \dots, q-1$, where $q \geq 2$, then I_q is an isomorphisme mod $\mathcal{C}$: This means that $\text{Ker}(I_q)$ and $\text{Coker}(I_q)$ are in $\mathcal{C}$.

The class of finitely generated Abelian groups is such an admissible class. In particular we have :

Theorem 4.2 *Let X be a simply connected space. Then $\pi_i(X)$ is finitely generated for $i \leq q$ iff $H_i(X)$ is finitely generated for $i \leq q$.*

In particular any closed, simply connected CW-complex has finitely generated homotopy groups (which is 1.1).

By applying this theorem to $\widetilde{M}$ we may replace the hypothesis on $\pi_i(M)$ by the analogue hypothesis on $H_i(\widetilde{M})$. From now on we will suppose that for all $i \leq \left\lfloor \frac{n}{2} \right\rfloor$, $H_i(\widetilde{M}) = \mathbf{Z}^{r_i} \oplus T_i$, where T_i is a torsion finitely generated $\mathbf{Z}$ -module. The proof of 4.1 relies on the following statement :

Proposition 4.3 *Let π be a group of type $\mathcal{F}_p$ for some positive integer p and $u : \pi \rightarrow \mathbf{Z}$ a morphism whose kernel is of type $\mathcal{F}_p$. Suppose that $\mathbf{Z}^r$ is a π -module. Denote by Λ the ring $\mathbf{Z}[\pi]$ and by Λ_u the completed ring, as above. Let $\pi_0 \leq \pi$ be a normal subgroup of finite index and Λ_0 the corresponding group ring $\mathbf{Z}[\pi_0]$. Then for $i \leq p$ we have*

$$\mathrm{Tor}_i^{\Lambda_0}(\mathbf{Z}^r, \Lambda_u) = 0.$$

Remark For $\pi_0 = \pi$ and $r = 1$ (and therefore for arbitrary r and trivial action of π on $\mathbf{Z}^r$) the statement above was proved by J-C. Sikorav in this thesis [32].

We postpone the proof of 4.3 to Subsection 4.4. We now show :

4.2 Proof of 4.3 $\implies$ 4.1

Fix $p, q \leq \left\lfloor \frac{n}{2} \right\rfloor$. For $g \in \pi_1(M)$, denote by ϕ_g the automorphism of $H_q(\widetilde{M}) = \mathbf{Z}^{r_q} \oplus T_q$, given by the action of $\pi_1(M)$. We have :

$$\phi_g = \begin{pmatrix} a_g & 0 \\ b_g & c_g \end{pmatrix},$$

where $a_g : \mathbf{Z}^{r_q} \rightarrow \mathbf{Z}^{r_q}$, $b_g : \mathbf{Z}^{r_q} \rightarrow T_q$ and $c_g : T_q \rightarrow T_q$. Note that a_g and c_g are automorphisms (of inverses $a_{g^{-1}}$, resp. $c_{g^{-1}}$). Let

$$\pi_0 = \{ g \in \pi_1 \mid c_g = Id \}.$$

Since T_q is finite, there is only a finite number of automorphisms $c : T_q \rightarrow T_q$. Therefore π_0 is a normal subgroup of $\pi_1(M)$ of finite index (It is the kernel of the morphism $\pi_1(M) \rightarrow \mathrm{Aut}(T_q)$ defined by $g \mapsto c_g$). We use the following :

Lemma 4.4 *Let G a group of type $\mathcal{F}_p$ and $G_0 \leq G$ a normal subgroup of finite index. Then G_0 is of type $\mathcal{F}_p$.*

The proof of 4.4 is obvious : If Q is a $K(G, 1)$ with finite p -skeleton, then the finite cover of Q corresponding to $G_0 \leq G$ will be a $K(G_0, 1)$ with finite p -skeleton.

Now let $u : \pi_1(M) \rightarrow \mathbf{R}$, as in the statement of 4.1 and let $u_0 = u|_{\pi_0}$. Obviously, $Ker(u_0)$ has finite index in $Ker(u)$ so, using 4.4, both π_0 and $Ker(u_0)$ are of type $\mathcal{F}_p$.

Consider now the short exact sequence :

$$0 \rightarrow T_q \xrightarrow{(0, Id)} \mathbf{Z}^{r_q} \oplus T_q \xrightarrow{Id \oplus 0} \mathbf{Z}^{r_q} \rightarrow 0.$$

One immediately checks that this is an exact sequence of π_0 -modules (where the action of π_0 on $\mathbf{Z}^{r_q}$ is $x \mapsto a_g(x)$). Now consider $\Lambda_0 = \mathbf{Z}[\pi_0]$, and view the completed ring Λ_u as a Λ_0 -module. The tensor product of Λ_u and the exact sequence above yields a long exact sequence :

$$(1) \quad \cdots \rightarrow Tor_p^{\Lambda_0}(T_q, \Lambda_u) \rightarrow Tor_p^{\Lambda_0}(\mathbf{Z}^{r_q} \oplus T_q, \Lambda_u) \rightarrow Tor_p^{\Lambda_0}(\mathbf{Z}^{r_q}, \Lambda_u) \rightarrow \cdots$$

As a consequence of 4.3, the right term in the sequence above vanishes. In order to prove that the left term is zero, we consider an exact sequence of the form

$$0 \rightarrow \mathbf{Z}^s \rightarrow \mathbf{Z}^m \rightarrow T_q \rightarrow 0,$$

which is viewed as a sequence of π_0 -modules with trivial actions (recall that, by construction, π_0 acts trivially on T_q). The corresponding long exact sequence given by the tensor product with Λ_u writes :

$$(2) \quad \cdots \rightarrow Tor_p^{\Lambda_0}(\mathbf{Z}^m, \Lambda_u) \rightarrow Tor_p^{\Lambda_0}(T_q, \Lambda_u) \rightarrow Tor_{p-1}^{\Lambda_0}(\mathbf{Z}^s, \Lambda_u) \rightarrow \cdots$$

Applying again 4.3 we infer $Tor_p^{\Lambda_0}(T_q, \Lambda_u) = 0$, therefore the middle term in (1) vanishes.

We have thus established that for each $p, q \leq \left\lfloor \frac{n}{2} \right\rfloor$

$$(3) \quad Tor_p^{\Lambda_0}(H_q(\widetilde{M}), \Lambda_u) = 0,$$

which is the assertion required in 4.1 ... with Λ_0 instead of Λ . To complete the proof we need the following results :

Proposition 4.5 *Let G be a groupe and let L be a $\mathbf{Z}[G]$ right module and N be a $\mathbf{Z}[G]$ left module. Assume that N is $\mathbf{Z}$ -torsion-free. Then*

$$Tor_*^{\mathbf{Z}[G]}(L, N) \approx H_*(G, L \otimes_{\mathbf{Z}} N),$$

where G acts diagonally on $L \otimes_{\mathbf{Z}} N : g(x \otimes y) = xg^{-1} \otimes gy$.

This proposition is proved in [7] (prop. 2.2, p. 61).

Theorem 4.6 *For any group extension*

$$1 \rightarrow K \rightarrow G \rightarrow Q \rightarrow 1$$

and any G -module R there is a spectral sequence of the form

$$E_{ij}^2 = H_i(Q, H_j(K, R))$$

which converges to $H_{i+j}(G, R)$.

This theorem is due to G. Hochschild and J-P. Serre [16] (see also [7], p. 171).

We apply first 4.5 and infer from (3) that for each $p, q \leq \left\lfloor \frac{n}{2} \right\rfloor$ we have :

$$(4) \quad H_p(\pi_0, H_q(\widetilde{M}) \otimes_{\mathbf{Z}} \Lambda_u) = 0.$$

Note that the hypothesis of 4.5 is fulfilled by $N = \Lambda_u$.

We fix q and denote by R the $\pi_1(M)$ -module $H_q(\widetilde{M}) \otimes_{\mathbf{Z}} \Lambda_u$ (for the diagonal action). Then we apply 4.6 to the extension

$$1 \rightarrow \pi_0 \rightarrow \pi_1(M) \rightarrow \pi_1(M)/\pi_0 \rightarrow 1$$

and to the module R ; We find using (4) that $E_{ij}^2 = 0$ for all $j \leq \left\lfloor \frac{n}{2} \right\rfloor$ and for all $i \in \mathbf{N}$. According to 4.6, this implies that

$$(5) \quad H_i(\pi_1(M), R) = 0 \quad \forall i \leq \left\lfloor \frac{n}{2} \right\rfloor.$$

Finally, we apply once again 4.5 and we get that for all $i, q \leq \left\lfloor \frac{n}{2} \right\rfloor$ we have :

$$Tor_i^{\Lambda}(H_q(\widetilde{M}), \Lambda_u) = 0.$$

The analogous relation for Λ_{-u} instead of Λ_u can be established in the same way, so 4.1 follows. $\diamond$

So we only have to proof 4.3 to complete the proof of 1.2, 1.3 and 1.4. The proof involves some facts about Bieri-Renz invariants. We recall the definition and some properties of these invariants in the subsection below :

4.3 Bieri-Renz invariants. Proof of 4.3

Let G be a group of type $\mathcal{F}_m$. We call two non zero homomorphisms $u, v : G \rightarrow \mathbf{R}$ equivalent if $u = \lambda v$ for some positive $\lambda \in \mathbf{R}$. We denote by $S(G)$ the quotient $\text{Hom}(G, \mathbf{R}) / \approx$ (which is a $(rk(G) - 1)$ -dimensional sphere). The Bieri-Renz invariants $\Sigma^i(G)$ and $\Sigma^i(G, \mathbf{Z})$, defined for $i = 1, \dots, m$ are open subsets of $S(G)$. They were introduced by R. Bieri, W Neuman and R. Strebel in [3] for $i = 1$ and by R. Bieri and B. Renz in [4] for $i \geq 2$.

These invariants are defined as follows. Let X be a $K(G, 1)$ which is a complex with finite m -skeleton and let $u : G \rightarrow \mathbf{R}$ be a non zero homomorphism. Then there exists an *equivariant height function* $f : \widetilde{X} \rightarrow \mathbf{R}$, i.e. a function which satisfies $f(gx) = f(x) + u(g)$. (If X is a manifold then f is a primitive of the pullback of some 1-form in the class u). It can be shown that the difference of two such functions is bounded. Consider the maximal subcomplex X_f of $\widetilde{X}$ whose image by f is contained in $[0, +\infty[$.

Definition. Let $u \in S(G)$ and $i \in \{1, \dots, m\}$. Then u belongs to $\Sigma^i(G)$ (resp. to $\Sigma^i(G, \mathbf{Z})$) if for some couple (X, f) as above the subcomplex X_f is $(i - 1)$ -connected (resp. $(i - 1)$ -acyclic).

Bieri and Renz show that the definition does not depend on (X, f) . The following properties of the Bieri-Renz invariants are obvious from the definition :

- a) $\Sigma^i(G) \subset \Sigma^i(G, \mathbf{Z})$.
- b) $\Sigma^1(G) = \Sigma^1(G, \mathbf{Z})$.
- c) For $i \geq 2$, $\Sigma^i(G) = \Sigma^i(G, \mathbf{Z}) \cap \Sigma^2(G)$.

The most striking application of these invariants is stated in the following :

Theorem 4.7 Let $N \subset G$ be a normal subgroup with Abelian quotient G/N . denote by $S(G, N)$ the subset of $S(G)$ defined by $\{u \in S(G) \mid u|_N = 0\}$. Then, for any $i = 1, \dots, m$, we have the equivalence :
 N is of type $\mathcal{F}_i$ iff $S(G, N) \subset \Sigma^i(G)$.

Note that $\{u, -u\} \subset S(G, \text{Ker}(u))$; When $\text{Im}(u)$ is cyclic it is easy to show that these two sets coincide. Note also that $S(G, [G, G]) = S(G)$. So, one immediately infers the following corollary :

Corollary 4.8 *i) Let $u \in S(G)$ and i an integer as above. If $\text{Ker}(u)$ is $\mathcal{F}_i$ then $\pm u \in \Sigma^i(G)$. The converse is valid if $\text{Im}(u)$ is cyclic.
ii) $\Sigma^i(G) = S(G)$ if and only if $[G, G]$ is of type $\mathcal{F}_i$.*

In the sequel we give an algebraic description of the invariants $\Sigma^i(G, \mathbf{Z})$ following [4], Section 4. Fix a non zero homomorphism $u : G \rightarrow \mathbf{R}$. Let F be a finitely generated free $\mathbf{Z}[G]$ -module, and $\{e_i\}_{i=1, \dots, k}$ a basis of F . We define an application $v : F \rightarrow \mathbf{R}$ as follows : v is defined arbitrarily on the elements e_i ; Then for any $g \in G$ we put $v(g e_i) = v(e_i) + u(g)$. Finally, if $\lambda \in F$ writes $\lambda = \sum_{i,j} n_{ij} g_j e_i$ in the basis $\{e_i\}$, we define

$$v(\lambda) = \inf\{v(g_j e_i) \mid n_{ij} \neq 0\},$$

and $v(0) = +\infty$. Following Bieri and Renz we call v a *valuation extending u* . Now suppose that G is of type $\mathcal{F}_m$ and let

$$P_m \xrightarrow{\partial_m} P_{m-1} \xrightarrow{\partial_{m-1}} \dots \xrightarrow{\partial_1} P_0 \xrightarrow{\partial_0} \mathbf{Z} \rightarrow 0$$

be a $\mathbf{Z}[G]$ -free, finitely generated resolution (which exists since there exists a $K(G, 1)$ with finite m -skeleton). Define as above $v_i : P_i \rightarrow \mathbf{R}$ which are valuations extending u . We may suppose in addition that for any $i = 1, \dots, m$ and for any $x \in P_i$ we have

$$(1) \quad v_i(x) \leq v_{i-1}(\partial_i(x)).$$

Indeed, one easily sees that it suffices to check the relation above on the basis $\{e_i\}$ of P_i . We can construct v inductively, by choosing $v(e_i)$ sufficiently negative in order to satisfy the inequality (1).

Bieri and Renz proved the following theorem ([4], theorem 4.1) :

Theorem 4.9 *Let $P_\bullet \rightarrow \mathbf{Z}$ be a $\mathbf{Z}[G]$ -free, finitely generated resolution of length m and let $v : P_\bullet \rightarrow \mathbf{R}$ be a valuation extending u satisfying (1). Then $u \in \Sigma^m(G, \mathbf{Z})$ if and only if there exists a chain endomorphism $\Phi : P_\bullet \rightarrow P_\bullet$ which lifts the identity of $\mathbf{Z}$ and which satisfies the property*

$$(2) \quad v(\Phi(x)) > v(x) \quad \forall x \in P_\bullet.$$

We will reformulate this theorem as follows. It is obvious that one can construct the valuation $v : P_\bullet \rightarrow \mathbf{R}$ such that it satisfies an additional feature : for all $j = 1, \dots, m$ v is constant on the set $\{e_i^j\}$ of the basis elements of P_j . Denote this constant by ν_j . Then, the valuation $v_j : P_j \rightarrow \mathbf{R}$ is given by

$$v_j(\lambda) = \nu_j + \inf\{u(g_k) \mid n_{ik}^j \neq 0\},$$

where $\lambda = \sum_{i,k} n_{ik}^j g_k e_i^j$, $g_k \in G$, $n_{ik}^j \in \mathbf{Z}$. Thus, for an endomorphism Φ given by 4.9, if

$$\Phi(e_i^j) = \sum_{i,k} n_{ik}^j g_{ik}^j e_k^j,$$

(where $g_{ik}^j \in G$ and $n_{ik}^j \in \mathbf{Z}$), the inequality (2) implies that $u(g_{ik}^j) > 0$ for all elements of G appearing with non zero coefficient n_{ik}^j .

We call an element $\lambda = \sum_i n_i g_i$ of $\mathbf{Z}[G]$ u -positive if $u(g_i) > 0$ for any g_i which has a non zero coefficient n_i in the writing of λ . We call a matrix $A \in \mathcal{M}_k(\mathbf{Z}[G])$ u -positive if all its entries are u -positive. Taking into account the preceeding remarks, theorem 4.9 can be stated as follows :

Theorem 4.10 *Let $P_\bullet \rightarrow \mathbf{Z}$ a resolution of length m which is $\mathbf{Z}[G]$ -free and finitely generated. Fix a basis for each P_j for all $j = 1, \dots, m$. Then $u \in \Sigma^m(G, \mathbf{Z})$ if and only if there exists a chain endomorphism $\Phi : P_\bullet \rightarrow P_\bullet$ which lifts the identity of $\mathbf{Z}$ and such that for all $j = 1, \dots, m$ the matrices of $\Phi_j : P_j \rightarrow P_j$ in the fixed basis are u -positive.*

Now we are able to complete the

Proof of 4.3

Applying 4.5 we infer that $\text{Tor}_i^{\Lambda_0}(\mathbf{Z}^r, \Lambda_u)$ is isomorphic to $H_i(\pi_0, \mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$, where the action of π_0 on $\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u$ is given by $x \otimes \lambda \mapsto xg^{-1} \otimes g\lambda$. It suffices therefore to prove that the latter vanishes for $i \leq p$.

Let

$$P_p \rightarrow P_{p-1} \rightarrow \dots \rightarrow P_1 \rightarrow P_0 \rightarrow \mathbf{Z}$$

be a $\mathbf{Z}[\pi_0]$ -free resolution which we may suppose finitely generated since π_0 is $\mathcal{F}_p$. By definition we have

$$(3) \quad H_i(\pi_0, \mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u) \approx H_i(P_\bullet \otimes_{\mathbf{Z}[\pi_0]} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)).$$

We will prove that the right term of (3) vanishes for all $i \leq p$. Fix a basis $\{e_i^j\}_i$ for each module P_j . Since $\text{Ker}(u|_{\pi_0})$ is of type $\mathcal{F}_p$, it follows by 4.8.i that $u \in \Sigma^p(\pi_0, \mathbf{Z})$, so, applying 4.10 we obtain an endomorphism $\Phi : P_\bullet \rightarrow P_\bullet$ such that for all $j = 1, \dots, m$ the matrix of Φ_j associated to the basis $\{e_i^j\}$ is u -positive.

Let $\Psi = \Phi \otimes \text{Id} : P_\bullet \otimes_{\mathbf{Z}[\pi_0]} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u) \rightarrow P_\bullet \otimes_{\mathbf{Z}[\pi_0]} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$.

The proof of 4.3 will be complete if we prove the following :

Lemma 4.11 *The homomorphism $\text{Id} - \Psi$ is invertible and it induces zero in homology.*

Proof

Let us prove first that $\text{Id} - \Psi$ vanishes in homology. Since $\Phi : P_\bullet \rightarrow P_\bullet$ lifts the identity of $\mathbf{Z}$, and $P_\bullet$ is free it is well known and easy to prove that there exists a homotopy $s : P_\bullet \rightarrow P_{\bullet+1}$ between Φ and Id . It follows that $s \otimes \text{Id}$ is a homotopy between Ψ and Id , so $\text{Id} - \Psi$ induces the zero morphism in homology.

Now let us prove the first assertion of the lemma. Let $\{f_1, f_2, \dots, f_r\}$ be the canonical basis of $\mathbf{Z}^r$. We can see $\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u$ as a right Λ_u -module endowed with the canonical structure : $(x \otimes \lambda)\mu = x \otimes \lambda\mu$. It is a free module of rank r and $\{f_1 \otimes 1, f_2 \otimes 1, \dots, f_r \otimes 1\}$ is a basis for this module.

Recall that we denoted by Λ_0 the ring $\mathbf{Z}[\pi_0]$. For any $j = 1, \dots, m$, the product $P_j \otimes_{\Lambda_0} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$ inherits of the structure of right Λ_u -module described above. On the other hand, since P_j is free we have

$$(4) \quad P_j \otimes_{\Lambda_0} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u) \approx (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)^{rk(P_j)} \approx (\Lambda_u)^{r \cdot rk(P_j)}.$$

If $\{e_i^j\}_{i=1, \dots, rk(P_j)}$ is the given basis of P_j then

$$(5) \quad \{e_i^j \otimes_{\Lambda_0} (f_s \otimes 1)\}_{i=1, \dots, rk(P_j), s=1, \dots, r}$$

will be a basis for $P_j \otimes_{\Lambda_0} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$.

It is easy to check that the isomorphisms (4) preserve the right Λ_u -module structure. Moreover, the differential $\partial \otimes_{\Lambda_0} \text{Id}$ of the complex $P_\bullet \otimes_{\Lambda_0} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$ respects this structure, therefore it is a complex free right Λ_u -modules.

We claim that the matrices of Ψ in the basis (5) are u -positive. Indeed, for fixed j , let $(\lambda_{ik})_{i,k=1, \dots, rk(P_j)}$ be the matrix of Φ_j in the basis $\{e_i^j\}$; This matrix is known as u -positive. We dropped down the index j from the coefficients

of the matrix to simplify the notations. Denote by $\bar{\lambda}$ the image of an element $\lambda \in \Lambda_0$ under the endomorphism $\Lambda_0 \rightarrow \Lambda_0$ induced by the involution of $\pi_0 : g \mapsto g^{-1}$. The right action of $\bar{\lambda}_{ik}$ on $\mathbf{Z}^r$, evaluated on the basis $\{f_s\}$ writes :

$$f_s \bar{\lambda}_{ik} = \sum_{l=1}^r n_{iks}^l f_l,$$

for some integers n_{iks}^l .

We infer that

$$\begin{aligned} \Psi_j(e_i^j \otimes_{\Lambda_0} (f_s \otimes 1)) &= \sum_k e_k^j \lambda_{ik} \otimes_{\Lambda_0} (f_s \otimes 1) = \\ &= \sum_k e_k^j \otimes_{\Lambda_0} (f_s \bar{\lambda}_{ik} \otimes \lambda_{ik}) = \sum_{k,l} e_k^j \otimes_{\Lambda_0} (n_{iks}^l f_l \otimes \lambda_{ik}) = \\ &= \sum_{k,l} e_k^j \otimes_{\Lambda_0} (f_l \otimes n_{iks}^l \lambda_{ik}) = \sum_{k,l} [e_k^j \otimes_{\Lambda_0} (f_l \otimes 1)] n_{iks}^l \lambda_{ik}, \end{aligned}$$

so the matrix of Ψ_j is u -positive.

Denote this matrix by A_j and define another matrix B_j by : $B_j = Id + \sum_{k=1}^{+\infty} A_j^k$. As A_j is u -positive it is easy to check that the matrix B_j belongs to $\mathcal{M}_{r \cdot rk(P_j)}(\Lambda_u)$. It therefore defines using the basis (5) an endomorphism of $P_j \otimes_{\Lambda_0} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$. It is actually an automorphism since obviously $(Id - A_j)B_j = Id$. Finally, as Ψ is a morphism of complexes, the morphism $Id + \Psi + \Psi^2 + \dots$ induced by B_j will also commute with the differential. We have finally got an automorphism of $P_{\bullet} \otimes_{\Lambda_0} (\mathbf{Z}^r \otimes_{\mathbf{Z}} \Lambda_u)$ whose inverse is $Id - \Psi$. This completes the proof of 4.11 and hence the proof of 4.3 which implies our main theorems 1.2, 1.3 and 1.4. $\diamond$

4.4 Proof of 1.5

Assume that the homotopy of X is finitely generated. By embedding X in $\mathbf{R}^{2q+3}$ construct a manifold $M = \partial W$ of dimension $2q+2$ as in the proof of 1.2. We have $\chi(X) = \chi(W) = 2\chi(M)$. By general position $\pi_i(M) \approx \pi_i(X)$ for $i \leq q+1$. These groups are therefore finitely generated. The hypothesis of 4.1 is fulfilled and, by applying this result, we get $H_*(M, u) = 0$. But according to 3.5 this implies $\chi(M) = 0$, contradicting thus the hypothesis on $\chi(X)$.

If X is a manifold, the result follows directly from 4.1, 3.1 and 3.5.

The proof is finished.

◇

Acknowledgements I am grateful to Thomas Delzant for our useful discussions on the subject.

Mihai DAMIAN
Université Louis Pasteur
IRMA, 7, rue René Descartes,
67 084 STRASBOURG

e-mail : damian@math.u-strasbg.fr

References

- [1] M. Bestvina, N. Brady, Morse theory and finiteness properties of groups, *Invent. Math.* 129 (1997), 445-470.
- [2] R. Bieri, B. Eckmann, Finiteness properties of duality groups, *Comment. Math. Helv.* 49 (1974), 74-83.
- [3] R. Bieri, W. Neuman, R. Strebel, A geometric invariant for discrete groups, *Invent. Math.* 129 (1987), 451-477.
- [4] R. Bieri, B. Renz, Valuations on free resolutions and higher geometric invariants of groups, *Comment. Math. Helv.* 63 (1998), 464-497.
- [5] G. Bredon, *Topology and Geometry*, GTM 139, Springer Verlag, New York, 1997.
- [6] W. Browder, J. Levine, Fiberings manifolds over the circle, *Comment. Math. Helv.* 40 (1966), 153-160.
- [7] K.S. Brown, *Cohomology of groups*, GTM 87, Springer Verlag, New York, 1982.
- [8] M. Damian, Formes fermées non singulières et propriétés de finitude des groupes *Ann.Sci.Ec.Norm.Sup.* 33 (2000) no.3, 301-320.
- [9] M. Damian, On the higher homotopy groups of a finite CW-complex, *Topology and its Appl.* 149 (2005), 273-284.

- [10] F.T. Farrell, The obstruction to fibering a manifold over the circle, *Bull. Amer. Math. Soc.* 73 (1967), 737-740.
- [11] F.T. Farrell, W.C. Hsiang, The Whitehead group of poly-(finite or cyclic) groups, *J. London Math. Soc.* (2) 24 (1981), 308-324.
- [12] K.W. Gruenberg, *Cohomological Topics in Group Theory*, L.N.M. 143, Springer Verlag, New York, 1970.
- [13] R. Godement, *Topologie algébrique et théorie des faisceaux*, Hermann, Paris, 1958.
- [14] K.A. Hirsch, On infinite soluble groups I, *Proc. London Math. Soc.* 44 (1938), 53-60.
- [15] K.A. Hirsch, On infinite soluble groups IV, *J. London Math. Soc.* 27 (1952), 81-85.
- [16] G. Hochschild, J.P. Serre, Cohomology of group extensions, *Trans. Amer. Math. Soc.* 74 (1953), 110-134.
- [17] B. Hu, Whitehead groups of finite polyhedra with non-positive curvature, *Jour. of Diff. Geom.* 38 3/1993, 501-517.
- [18] F. Latour, Existence de 1-formes fermées non-singulières dans une classe de cohomologie de de Rham, *Publ. Math. IHES* 80 (1994).
- [19] S. Maumary, Type simple d'homotopie, L.N.M. 48, Springer Verlag, New York, 1967.
- [20] J. Meier, H. Meinert, L. Van Wyk, Finiteness properties and abelian quotients of graph groups, *J. Math. Res. Lett.* 3 (1996), 779-785.
- [21] J. Milnor, Whitehead torsion, *Bull. Amer. Math. Soc.* 72 (1966), 358-426.
- [22] S.P. Novikov, Multivalued functions and functionals. An analogue of the Morse theory, *Soviet. Math. Dokl.* vol 24, nr. 2/1981, 222-226.
- [23] A. Pajitnov, Surgery on the Novikov complex, *K-theory* 10 (1996), 323-412.

- [24] A. Pajitnov Surgery on the Novikov complex, Rapport de Recherche, Nantes (1993), <http://193.52.98.6~pajitnov>.
- [25] A. Pajitnov, A. Ranicki, The Whitehead group of the Novikov ring, *K-theory* 21 (2000) no.4, 325-365.
- [26] A. Ranicki, High-dimensional knot theory, Springer (1998).
- [27] A. Ranicki, Lower K- and L- theory, L.M.S. Lecture Notes, Cambridge, 1992.
- [28] A. Ranicki, Finite domination and Novikov rings, *Topology* 34 (1995), 619-632.
- [29] J. Stallings, The piecewise linear structure of the Euclidean space, *Proc. Cambridge Phil. Soc.* 58 (1962), 481-488.
- [30] J.P. Serre, Groupes d'homotopie et classes de groupes abéliens, *Ann. of Math.* (2) 58 (1953), 258-294.
- [31] L. Siebenmann, A total Whitehead obstruction to fibering over the circle, *Comment. Math. Helv.* 45 (1970), 1-48.
- [32] J-C. Sikorav, Homologie de Novikov associée à une classe de cohomologie réelle de degré un, Thèse Orsay 1987.
- [33] E. Spanier, Algebraic Topology, Mc Graw-Hill, New York, 1966.
- [34] D. Tischler, On fibering certain foliated manifolds over S^1 , *Topology* 9 (1970), 153-154.
- [35] C.T.C. Wall, Finiteness conditions for CW-complexes, *Ann. of Math.* 81 (1965), 56-69.